Fig. 1. G-CSF Synergizes IL-8 Induced Neutrophil Chemotaxis




Figure 2 GM-CSF Synergizes IL8 Induced PMN Chemotaxis





Fig. 3. Dose Response Curve for IL-8 with Constant G-CSF (100 pM)



IL-8 (nM)

Fig. 4. GCSF Does not Synergize f-MLP Induced Neutrophil Chemotaxis





Fig.5. G-CSF enhances in vivo neutrophil intradermal recruitment



Minus baseline & normalized against tissue weight

Fig. 6. Binding of ¹²⁵I G-CSF on PMN




Fig. 7. G-CSF Neutralizing Antibody Inhibits G-CSF Synergized Chemotaxis





Fig. 8. G-CSF Pre-Incubation Decreases Neutrophil Response to IL-8



Cells were preincubated with G-CSF for respective time periods and subsequently treated with 1nM of IL-8

Fig. 9. G-CSF Does not Alter IL-8 Induced Calcium Flux






Fig. 10. G-CSF Does Not Increase IL-8 Binding in Neutrophils



Scintillation Count (x Thousands)

D988889,022801

Fig. 11. G-CSF Preincubation Does not Alter IL-8 Binding on Neutrophils



100 pM of G-CSF was incubated simultaneously or pretreated for the respective time periods

Fig. 12 G-CSF Pre-Incubation Alters PMN Response to LI-8




Figure 13: G-CSF potentiates both chemokinetic and chemotactic effects of IL-8




Figure 14: FACS dot plot of FSC vs. SSC from unstimulated human whole blood




Figure 15: FACS Dot plot of FSC vs. SSC from unstimulated and MCP-1 stimulated human whole blood






Figure 16: Time course of FSC changes in response to MCP-1 stimulation S FSC (% of control)





M-CSF, pM

Figure 19: Effect of M-CSF on human monocyte shape change - comparison to MCP-1

